

MISOCLEAR

Misoprostol (200mcg)

High quality, effective, proven

For early abortion of pregnancy up to 9 weeks

































Misoprostol Dosage by Indication and Gestation

*Misoclear should only be used for approved indications permitted by respective national regulation.

Please see
reverse for
guide to
pictures
showing route
of dosage

General warning

Misoprostol is a very powerful stimulator of uterine contractions in late pregnancy and can cause fetal death and uterine rupture if used in high doses. Please use clinical judgement in dosing.

Indication	Instructions for Each Dose				Notes
Cervical priming pre-instrumentation	 400mcg		Vaginally 3-4 hours before procedure Repeat in 3-4 hours if necessary	OR  Sublingually 2-3 hours before procedure	
Induced abortion (up to 12 weeks)	 800mcg		Vaginally every 3 hours Max. 3 doses	OR  Sublingually every 3 hours Max. 3 doses	Third dose only taken if pregnancy has not been passed
Induced abortion (>12-24 weeks)	 400mcg		Vaginally every 3 hours Max. 5 doses	OR  Sublingually every 3 hours Max. 5 doses	
Treatment of Incomplete Abortion (up to 12 weeks) [To stop post-abortion bleeding due to retained products]	 600mcg OR  400mcg		Orally as a single dose		Sublingually as a single dose
Treatment of Incomplete Abortion (>12-24 weeks)	 400mcg		Vaginally every 3 hours Max. 5 doses	OR 	Sublingually every 3 hours Max. 5 doses
Treatment of Missed Abortion (Up to 12 weeks) [Pregnancy has terminated but no products of conception expelled]	 800mcg OR  600mcg		Vaginally every 3 hours Max. 3 doses		Sublingually every 3 hours Max. 3 doses
Treatment of Intra-Uterine Fetal Death (>12 up to 17 weeks)	 200mcg		Vaginally every 6 hours Max. 4 doses		Halve doses in women with previous caesarean section or uterine scar
Treatment of Intra-Uterine Fetal Death (>17 up to 26 weeks)	 100mcg (half a tablet)		Vaginally every 6 hours Max. 4 doses		
Treatment of Intra-Uterine Fetal Death (>26 weeks)	 25mcg dissolved in solution		25mcg dose taken orally in dissolved solution* Max. 8 doses		*Please see oral dissolved solution instructions on next page
AND Induction of labour	 25mcg pill if available		Vaginally every 6 hours Max. 4 doses		Do not use if previous caesarean section or uterine scar
PPH Prophylaxis (Prevention)	 600mcg		Orally as a single dose		
PPH treatment	 800mcg		Sublingually as a single dose		

MISOCLEAR

Misoprostol (200mcg)

High quality, effective, proven
For early abortion of pregnancy up to 9 weeks

Guide to Dosage Routes

Route	Instructions for Use
<div>Oral</div> <div></div>	<p>Pills are swallowed</p>
<div>Sublingual</div> <div></div>	<p>Pills are placed under the tongue, held for 30 minutes and then swallowed</p>
<div>Vaginal</div> <div></div>	<p>Pills are inserted in the Vaginal Fornices (deepest portions of the vagina) and the woman then lies down for 30 minutes</p>
<div>Oral Dissolved Solution</div> <div></div>	<p>One 200mcg pill is dissolved in 200mL of water to make a 200mcg/200mL solution. Client is then given 25 mL, which is equal to 25mg</p> <p>Note that breaking Misoclear tablets into smaller pieces than halves is not recommended. If a 25mcg strength pill is available, this may be used vaginally as shown in the table</p>